

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## GEOMETRY.

## 338. Proposed by C. N. SCHMALL, 239 East 7th Street, New York.


Given the base and vertical angle of a triangle, find the locus of the center of its nine-point circle.

### I. Solution by J. W. CLAWSON, Ursinus College, Collegeville, Pa.

(1) Take an angle BAC equal to the given vertical angle and from any point B in one arm of it describe an arc with BC equal to the given base as

radius cutting the other arm in C. Draw a circle through A, B, C. Then BC being fixed, the circle ABC is the locus of the vertex A.

(2) Find the center O of this circle, draw OA' the perpendicular bisector of BC, and OB' the perpendicular bisector of CA. Drop AD perpendicular to BC, and BE perpendicular to CA, intersecting AD in H.



The nine-point circle passes through A', D and B', E. Therefore the lines bisecting A'D and B'E at

right angles intersect in N, the center of the nine-point circle. But evidently each of these lines bisects OH. Therefore N bisects OH.

- (3) Join A'B'. The triangles AHB and OA'B' are similar, having their sides mutually parallel. Also AB=2A'B'. So AH=2OA', and is therefore constant.
- (4) Since AH is constant in length, and as A moves AH moves parallel to itself, the locus of H is a circle of radius equal to OA' and whose center, Q, lies on OA' produced so that OQ=AH=2OA'. Draw this circle.
- (5) Since N bisects OH, O is a fixed point, and the locus of H is a circle, the locus of N is a circle, whose center bisects OQ, and is therefore the point A', and whose radius is half the radius of the original circle.

Also solved by J. M. Meyer, S. J., Francis Rust, V. M. Spunar, G. B. M. Zerr, and J. Scheffer.

# CALCULUS.

#### 265. Proposed by V. M. SPUNAR, M. and E. E., East Pittsburg, Pa.

Find two curves which possess the property that the tangents TP and TQ to the inner one always make equal angles with the tangent TT' to the outer.

#### II. Solution by F. H. SAFFORD, Ph. D., University of Pennsylvania.

Let  $y=\phi(x)$  be the equation of the inner curve, and let the coordinates of P and Q, points on this curve, be, respectively,  $(x_1, y_1)$ ,  $(x_2, y_2)$ . Assuming that the outer curve exists, let its equation be y=f(x), and let  $f'(x)=\tan \alpha$ .

The two tangent lines from P and Q, respectively, are, if  $\phi'(x) = \tan \theta$ ,